

Instruction of Nitrostat (Nitroglycerin) Sublingual Tablet

1. Nitrostat[®] sublingual (under the tongue) tablet is used to treat or prevent angina (chest pain). This drug is taken on an as needed basis.
2. (1) Sit down before use. This medicine may make you dizzy. After using medication, you may feel lightheaded when standing, so stand up slowly. (2) Place a Nitrostat[®] sublingual tablet under your tongue and let melt. (3) Additional sublingual tablets may be used at 5-minute intervals, if chest pain is not relieved. (4) If chest pain persists after a total dose of 3 tablets in a 15-minute period, call the Emergency Department or get a ride to the hospital. Do not drive yourself.
3. Store the medicine at room temperature (20 ~ 25°C) in the original container, tightly capped, away from heat, moisture, and direct light. Nitrostat[®] shall be dated when first opened and expire 6 months after the opening date.
4. Nitrostat[®] is a nitrate. You should not use this medicine if you are also using Viagra[®], Cialis[®], or Levitra[®], since these drugs may potentiate the hypotensive effects of organic nitrates. Check all drugs you are taking with your doctor.
5. Other side effects: headache, facial flushing, nausea, vomiting, weakness. Alcohol can worsen the side effects.

TAH Drug Identity Search System

