



Instruction of Oral Antiviral Suspension for Influenza

● Generic Name: Oseltamivir (Treatment of influenza)

Trade Name	Chinese Name	TAH Drug Identify Search System
Tamiflu 75mg	克流感膠囊	
Eraflu 75mg	易剋冒膠囊	

● Preparation: The following are two examples of different dosage.

Dosage/each time	Instructions
0.27 Capsule (20 g)	<ol style="list-style-type: none"> 1. Open three capsules, then add the powder of the capsules into a small bottle. 2. Shake the bottle to loosen the powder. 3. Add cold water below the line or label. Invert the bottle and shake well, then top up with water exactly to the line (45 mL). Invert the bottle and again shake well. 4. Once reconstituted, store in a refrigerator (2-8°C, but not frozen). 5. Each time take 4 mL, twice a day, in the morning and evening. 6. Shake the oral suspension before use.
0.16 Capsule (12 mg)	<ol style="list-style-type: none"> 1. Open two capsules, then add the powder of the capsule into a small bottle. 2. Shake the bottle to loosen the powder. 3. Add cold water below the line or label. Invert the bottle and shake well, then top up with water exactly to the line (30 mL). Invert the bottle and again shake well. 4. Once reconstituted, store in a refrigerator (2-8°C, but not frozen). 5. Each time take 2.4 mL, twice a day, in the morning and evening. 6. Shake the oral suspension before use.



● Precaution



1. Follow doctor's advice and use within 5 days.
2. Avoid missing doses and take this medicine at the same time every day, in the morning and evening.
3. Tamiflu® and Eraflu® are available as capsules containing 75 mg oseltamivir, prepare concentration of oral antiviral suspension is 5mg/mL.

Instruction of Oral Antiviral Suspension for Influenza

● Generic Name: Oseltamivir (Treatment of influenza)

Trade Name	Chinese Name	TAH Drug Identify Search System
Tamiflu 75mg	克流感膠囊	
Eraflu 75mg	易剋冒膠囊	

● Preparation: Please Prepare according to the following table

Dosage/each time	Instructions
<p>_____ Capsule (_____ mg)</p>	<ol style="list-style-type: none"> 1. Open _____ capsules, then add the powder of the capsule into a small bottle. 2. Shake the bottle to loosen the powder. 3. Add cold water below the line or label. Invert the bottle and shake well, then top up with water exactly to the line (_____ mL). Invert the bottle and again shake well. 4. Once reconstituted, store in a refrigerator (2-8°C, but not frozen). 5. Each time take _____ mL, twice a day, in the morning and evening. Shake the oral suspension before use.

● Precaution

1. Follow doctor's advice and use within 5 days.



2. Avoid missing doses and take this medicine at the same time every day, in the morning and evening.
3. Tamiflu® and Eraflu® are available as capsules containing 75 mg oseltamivir, prepare concentration of oral antiviral suspension is _____mg/mL.